

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323831888>

Patellofemoral pain syndrome in female athletes: A review of diagnoses, etiology and treatment options

Article in *Orthopedic Reviews* · March 2018

DOI: 10.4081/or.2017.7281

CITATIONS

0

READS

73

5 authors, including:



Emily Curry

Boston University

32 PUBLICATIONS **125** CITATIONS

[SEE PROFILE](#)



Elizabeth Matzkin

Brigham and Women's Hospital

42 PUBLICATIONS **604** CITATIONS

[SEE PROFILE](#)



Xinning Li

Boston University

104 PUBLICATIONS **867** CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Commentary & Perspective Total Hip Arthroplasty [View project](#)

Patellofemoral pain syndrome in female athletes: A review of diagnoses, etiology and treatment options

Molly Vora,¹ Emily Curry,²
Amanda Chipman,³ Elizabeth Matzkin,⁴
Xinning Li¹

¹Boston University School of Medicine, MA; ²Boston University School of Public Health, MA; ³Tufts University School of Medicine, Boston, MA; ⁴Harvard Medical School, Boston, MA, USA

Abstract

Patellofemoral pain syndrome (PFPS) is one of the most common causes of knee pain and is present in females disproportionately more relative to males. PFPS causes tend to be multifactorial in nature and are described in this review. From a review of the current literature, it is clear that there needs to be further research on PFPS in order to better understand the complex etiology of this disorder in both males and females. It is known that females with patellofemoral pain syndrome demonstrate a decrease in abduction, external rotation and extension strength of the affected side compared with healthy patients. Conservative management, including optimizing muscle balance between the vastus medialis and lateralis around the patella along with formal therapy should be the first line of treatment in patients presenting with PFPS. Surgery should be reserved for patients in which all conservative management options have failed. This review aims to guide physicians in accurate clinical-decision making regarding conservative and surgical treatment options when specifically faced with PFPS in a female athlete. Furthermore, we will discuss the anatomic variants, incidence and prevalence, etiology, diagnosis and treatment of PFPS.

Introduction

Patellofemoral pain syndrome (PFPS) is the most common cause of knee pain in female athletes and is a result of imbalances in the forces controlling patellar tracking during knee flexion and extension (Table 1).¹ Symptoms include pain behind or around the patella that is increased with running or other knee flexion activities, such as squatting and walking up and down stairs. PFPS is disproportionately present in

females relative to males.² The goal of this review is to guide physicians in making accurate clinical decisions when evaluating PFPS in a female athlete.

Knee anatomy and patellofemoral pain syndrome

The patellofemoral joint consists of the patella, the distal and anterior aspects of the femur as well as the articular surfaces and surrounding supporting structures.³⁻⁵ The patella is the largest sesamoid bone in the body and is of a relatively constant length, width and thickness.^{6,7} Seventy-five percent of the posterior aspect of the patella is covered by cartilage up to five millimeters thick. This cartilage has both elastic and viscous properties. The fluid component allows for force absorption and lubrication of the articular surface, while the elastic portion helps to distribute and absorb forces.⁸⁻¹⁰ The tendons of the four components of the quadriceps muscle converge in the distal portion of the thigh and unite to form a single broad quadriceps tendon. The patellar tendon, which inserts on the tibial tuberosity is the continuation of this quadriceps tendon in which the patella is embedded. The medial and lateral vasti muscles of the quadriceps also attach independently to the patella and form aponeuroses, known as the medial and lateral patellar retinacula, respectively.^{7,11}

PFPS is the name given to a variety of pathologies that lead to anterior knee pain. PFPS is difficult to define because patients experience a variety of symptoms and may have different levels of pain and physical impairment.^{8,12} Further, most current literature focuses on studies performed with male participants, limiting the knowledge of treatment options for females with PFPS.

Incidence and prevalence

PFPS is the most prevalent orthopedic condition seen in sports medicine and is a common presenting complaint in adolescents and young adults.^{13,14} PFPS is also the primary diagnosis in about 25% of all running injuries.^{5,15} Treatment for PFPS is especially promising for the short term, but long-term results are much less successful.¹⁶ The total incidence for PFPS ranges from 8.75% to 17%; however, the incidence among females is much greater at 12.7% compared to 1.1% of males.¹⁷ Young females who regularly participate in running and jumping activities may be particularly at risk.² In a clinical analysis of 40 women with PFPS, pain was

Correspondence: Xinning Li, Department of Orthopedic Surgery, Sports Medicine and Shoulder Surgery, Boston University School of Medicine - Boston Medical Center, 850 Harrison Avenue - Dowling 2 North, Boston, MA 02118, USA.
Tel.: +1.508.816.3939.
E-mail: xinning.li@gmail.com

Key words: patellofemoral pain syndrome, PFPS, anterior knee pain, female athletes, treatment options

Contributions: MV, made substantial contributions to the acquisition of data, formatting of the manuscript and drafting the manuscript. EJC, made substantial contributions to the conception and design of the work and interpretation of the data for the work. AC, made substantial contributions to the conception of the work. She also was responsible for the acquisition of the data for the work and drafting the work. EM, made substantial contributions to the conception of the work and interpretation of the data for the work. She also was involved with the critical revision for important intellectual content. XL, made substantial contributions to the conception of the work and interpretation of the data for the work. He also was involved with the critical revision for important intellectual content.

Conflict of interest: MV, EJC, AC, declares no conflicts of interest relevant to this submission. EM, declares no conflicts of interest relevant to this submission, but receives research support from Zimmer. XL declares no conflicts of interest relevant to this submission, but is on the editorial board of JoMI and also holds equity in the company (<\$5,000).

Received for publication: 25 June 2017.
Revision received: 8 October 2017.
Accepted for publication: 8 October 2017.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright M. Vora et al., 2017
Licensee PAGEPress, Italy
Orthopedic Reviews 2017;9:7281
doi:10.4081/or.2017.7281

associated with increased activity. Chronic overloading and overuse of the patellofemoral joint, rather than malalignment, can also contribute to patellofemoral pain.¹⁸ A study of freshmen at the United States Naval Academy conducted by Boling *et al.*¹⁹ found that females were 2.23 times more likely to develop PFPS compared with males. Additionally, Boling *et al.* found that the prevalence of PFPS was not significantly different between sexes at the time of admission to the US Naval Academy. This data

along with other studies suggests that females are more affected than males by a rapid increase in physical activity level, which in turn leads to a higher incidence of PFPS.^{19,20}

Etiology

The causes of PFPS in females are multifactorial and include overuse injuries of the extensor apparatus (tendonitis or insertional tendinosis), patellar instability, chondral and osteochondral damage.²¹

Malalignment of the lower extremity

Malalignment of the lower extremity has been cited as a potential contributory factor in the development of PFPS. Femoral neck anteversion, genu valgum, knee hyperextension, Q angle, tibia varum and excessive rearfoot pronation are some of the alignment factors that have been associated with PFPS.⁸

Q angle is defined as the angle between the line connecting the anterior superior iliac spine to the center of the patella and the extension of a line from the center of the patella to the tibial tubercle.²¹ A greater lateralization angle is exerted on the patella with a greater Q angle, which increases the load on the lateral facet of the patella and the lateral femoral condyle. A 10% increase in the Q angle will result in increased stress to the patellofemoral joint by 45%.²² A Q-angle greater than 20 degrees for women is considered clinically abnormal.²³ While some data suggests that a greater Q angle is not a risk factor for PFPS, others suggest that a high Q angle may be a contributing factor in maintaining PFPS once it has been acquired.²⁴ Additionally, some authors have attributed excessive dynamic knee valgus malalignment in patients with PFPS compared to normal patients.²⁵

Patterns of patellar malalignment include subluxation with and without patellar tilt as well as patellar tilt without subluxation.⁸ In a computerized tomography study of the patellofemoral joint during active flexion and extension, lateral patellar translation and tilt was present in 8 out of 20 knees with anterior knee pain.²⁶ Abnormal surface tracking at the patellofemoral joint has often been cited as a potential cause of PFPS.²⁷ Patellar tracking, which targets dynamic patellofemoral alignment throughout knee range of motion, is essential for healthy joint function and affects contact and load transmission.²⁸ Cartilage thickness also has been suggested to influence joint contact and as a result may be another contributing factor to PFPS.^{29,30} Furthermore, in patients with trochlear dysplasia and

patellofemoral instability, Shin *et al.*³¹ reported decreased trochlear volume and length compared to normal control groups. Thus, normal patellofemoral tracking is dependent on many factors.

Although dynamic lateral patella maltracking is a risk factor for PFPS, static patellar malalignment can also be a contributing factor.³² Differential action of the quadriceps, in particular the vastus medialis obliquus, has been involved in the etiology of PFPS.³³ Lin *et al.*³² observed that vastus medialis obliquus activation in PFPS patients caused greater medial patellar rotation than in healthy subjects. Additionally, they reported that the three-dimensional kinematic action of the vastus medialis obliquus is actively modulated with knee flexion angle in healthy subjects, but that this modulation was not present in PFPS patients. These results could be attributed to differential vastus medialis obliquus insertion on the patella or medial quadriceps weakness.

Muscular imbalances

Decreased strength due to atrophy or inhibition of the lower extremity musculature has been suggested as a possible cause for PFPS.⁸ There are a number of muscular imbalances that are thought to contribute to PFPS development and include decreased knee extensor strength, weakness in eccentric muscle strength, imbalance between the vastus medialis obliquus and vastus lateralis components of the quadriceps, and hip muscle weakness.³⁴ Studies have shown that quadriceps atrophy is associated with PFPS pain syndrome.³⁵⁻³⁷ However, Thijs *et al.*³⁴ observed that the strength of hip muscle groups in female runners who developed patellofemoral pain did not significantly differ from those of the asymptomatic runners.³⁴ Other more recent studies suggest that female athletes with greater hip abduction strength might be at an increased risk of developing PFPS.³⁸

Decreased knee extensor strength: quadriceps volume and strength deficiency

Decreased knee extensor strength is a common finding in patients with PFPS.⁸ Thomee *et al.*⁸ found that patients with PFPS have more symptoms and pain during the last thirty degrees of maximal sitting extension. A study on young women with PFPS showed significantly lower knee extensor strength in the symptomatic knee. Further, the patients had less vertical jumping ability and were weaker, with the largest differences in eccentric knee extension. Affected patients had lower strength, EMG activity and significant differences in mus-

cle activity between the vastus medialis and rectus femoris muscle. Another feature of PFPS is decreased knee extensor torque. Kaya *et al.*³⁹ showed that women with PFPS have a decreased torque, total volume, and cross sectional area of the quadriceps muscle. Decreased torque also leads to muscular imbalances that increases the risk of PFPS.

Lephart *et al.*⁴⁰ indicated that females have significantly more hip internal rotation to maximum angular displacement, and less lower leg internal rotation time to maximum angular displacement compared to males. Females also have significantly less peak torque to body mass for the quadriceps and hamstrings than males. Weaker thigh musculature could be associated with stiffening of the knee and lower leg upon landing in females.⁹ Additionally, Besier *et al.*⁴¹ reported that PFPS patients had greater contraction of quadriceps as well as hamstrings and greater normalized muscle forces during walking, although the net knee moment was similar between PFPS patients and healthy pain-free controls. Females displayed 30-50% greater normalized gastrocnemius and hamstring muscle forces during both running and walking when compared to males.

Vastus medialis obliquus insufficiency and atrophy

Vastus medialis obliquus imbalance relative to the vastus lateralis has been cited as one of the main contributors to abnormal patellar tracking.³⁴ Under normal conditions, the vastus medialis obliquus and vastus lateralis counteract each other and are considered to be important patellofemoral joint stabilizers.⁴² When the balance of the vastus medialis obliquus and vastus lateralis is disrupted, it is often attributed to insufficiency of the vastus medialis obliquus due to atrophy, hypoplasia, inhibition or impaired motor control.⁴³ Hence it has been suggested that PFPS is linked to a decrease in vastus medialis obliquus muscle mass.^{34,44} The insertion of the vastus medialis obliquus is along the medial border of the patella and it extends from one third to one half of the way down from the proximal pole. Jan *et al.*⁴⁵ found that insertion level of the vastus medialis obliquus was significantly higher in patients with PFPS than healthy controls. Further the vastus medialis obliquus fiber angle was significantly smaller than in healthy control knees.

Differential activation of vastus medialis obliquus versus vastus lateralis

Another theory regarding the etiology of PFPS suggests that there is a differential activation time between vastus medialis

obliquus and vastus lateralis. Poor coordination of activation onset times of vastus lateralis and vastus medialis obliquus can lead to abnormal patella tracking.⁴⁶ Akkurt *et al.*⁴⁷ reported a significant delay in the activation onset time of vastus medialis obliquus in the affected knee of female patients at fifteen, thirty and forty-five degree knee extension angles as measured by electromyographic recording. They also reported that the delay in female patients was more pronounced at knee angles closer to full extension. Conversely, Karst *et al.*⁴⁸ found no difference in the initial activation of vastus medialis obliquus and vastus lateralis activities in patients with PFPS and asymptomatic individuals during three testing conditions: reflex knee extension, active knee extension in non-weight bearing and weight bearing situations. In a systematic review and meta-analysis of the literature, Chester *et al.*⁴⁹ evaluated 14 studies comparing the timing of EMG onset of VMO and VL in patients with PFPS versus asymptomatic individuals and found considerable heterogeneity between each study design. Although the data indicate a trend towards a delay in the VMO activation relative to VL in the PFPS patient population during both the voluntary task and reflex activities, the authors could not draw a clinical or therapeutic significance due to the variability in physiological function among normal individuals.

Hip muscle weakness

While hip muscle weakness is not directly associated with the patellofemoral joint, it is often associated with PFPS. The kinetic chain theory states that dysfunction of a joint can manifest injuries in other joints, most usually those distal to the affected joint.³⁴ It has been demonstrated that during running, females exhibit significantly greater external knee valgus movement and hip internal rotation than males.⁴⁰ The ability to control and prevent these motions relies on the strength of the proximal muscle groups that are antagonistic to these movements. If there is not sufficient proximal strength, the femur may adduct or internally rotate, which in turn increases lateral patellar contact pressure which may lead to pain.⁵⁰ Ireland *et al.*⁵⁰ reported that female PFPS patients had 26% less hip abduction strength and 36% less hip external rotation strength. Other similar studies conducted in a sample of females reported results that were in agreement with this study.^{51,52}

Gluteal muscle activation

It has been shown that altered hip joint kinematics is demonstrated in patients with

PFPS. Females with PFPS have demonstrated increased hip internal rotation during single step downs, running and jumping.⁵³ Increased hip adduction and knee abduction during walking have also been observed in female patients with PFPS. These transverse and frontal plane rotations are thought to reduce patellofemoral contact area and increase patellofemoral joint stress, which leads to pain. It has been proposed that altered neuromuscular control of the musculature that resists hip adduction and internal rotation may contribute to the kinematic differences observed between females with PFPS and healthy controls.⁵³

Overuse

Although many studies have attempted to explain the etiology of PFPS, most have focused on muscular imbalances and biomechanical abnormalities. However, physical activity level and overuse is also an important factor in the development of PFPS.⁸ Fairbank *et al.*⁵⁴ reported that female patients with PFPS were more involved in competitive sports than age-matched controls and that pain was related to increases in physical activity level. In fact, Thomee *et al.*¹⁸ found that all female patients who reported symptoms of an insidious onset of PFPS had been involved in temporary overuse or a period of increased physical activity. Interestingly, females with a high physical activity level did not report more pain than those with a lower activity level.¹⁸ This suggests that a drastic increase or change in activity is the stimulus leading to PFPS development rather than a consistently high level of activity.

Clinical evaluation

Females (62% of cases) are at a significantly greater risk of experiencing patellofemoral pain syndrome than males (38% of cases).⁵⁵ Anatomic, hormonal factors and knee laxity, and neuromuscular factors contribute to the higher risk, with anatomic factors being the most commonly discussed.^{56,57} One of the neuromuscular factors lacking in females is hip muscle strength.⁵⁸ Further, females have less hip external rotation and abduction strength than men. Female athletes who suffered a lower extremity injury during the season had a significant deficit in hip abduction and external rotation strength compared to injured controls.⁵⁹ Excessive femoral adduction and internal rotation may increase the dynamic quadriceps angle and lead to greater lateral patellar contact pressure.²⁸ In order to evaluate PFPS, the clinician

must first evaluate the lower extremity alignment and the extensor mechanism.²⁰ The clinician should aim to distinguish between an alignment problem within the patellofemoral joint, an alignment issue outside of the patellofemoral joint or absence of malalignment.²⁰ Within the so called miserable malalignment syndrome, a CT Scan is necessary to diagnose torsion, rotation and femoral neck anteversion correctly.^{60,61} The first step is visual inspection of the lower extremity with the feet together. Full-length alignment radiographs is essential for the accurate diagnosis and measurement of malalignment to guide management.

The source of pain in PFPS may not always be malalignment or patella instability, but instead excessive loading of the patellofemoral joint. The excessive loading may be a consequence of a single event or may be chronic in nature.⁶² In the majority of PFPS patients, no abnormal anatomical or biomechanical reasons for the symptoms exist.²⁰ Both a static and dynamic evaluation of the entire leg should be performed. The patient should stand and walk barefoot while the alignment and functionality of the lower extremity is evaluated. If there is a functional abnormality, the clinician should determine the reason for this compensatory mechanism, such as muscle weakness, muscle tightness or patellar hypermobility. The clinician should also evaluate for discrepancies in leg length and intrinsic foot imbalances. If an intrinsic foot imbalance exists, orthotics may be included as part of an effective treatment program.⁶³ If malalignment of the lower extremity is observed, mobilizing techniques and a formal exercise and stretching program can be used to correct postural and movement dysfunction.²⁰ Additionally, the clinician can incorporate 5 functional performance tests which include anteromedial lunge, step-down, single-leg press, bilateral squat, balance and reach to further assess patient progress.⁶⁴

The patella should be evaluated for glide, medial and lateral tilt, anterior and posterior tilt, and rotation. Exam findings should be compared to the contralateral normal side as some patients have excessive laxity but without any pain. Examination using radiographs, in particular the sunrise view to evaluate patella tilt and CT should be used for further evaluation of patellar tracking. MRI should be utilized to rule out meniscal, ligamentous or cartilage pathology. The condition of the non-muscular tissue around the patellofemoral joint must also be examined. The medio-lateral displacement test allows the clinician to reliably test the mobility of the patella and to determine whether it is normal, hypermobile or hypomobile. Previous studies show a

possible link between the role of the menstrual phase and hormonal factors in the development of PFPS. An increase in knee laxity and other ligament mechanical properties caused by fluctuations in female sex hormones may increase the risk of ACL injuries and PFPS.⁶⁵ Tenan *et al.*⁵⁷ showed that the vastus medialis and vastus medialis oblique initial firing rates vary throughout the menstrual cycle.

Sinding Larsen's disease, Osgood Schlatter's disease, and neuromas.

Patellar subluxation, dislocation, or prior surgery may lead to articular cartilage injury which also results in anterior knee pain. Risk factors for PFPS in females include overuse, trauma, muscle dysfunction, tight lateral restraints, patellar hypermobility, and poor quadriceps flexibility (Table 2).

developing rehabilitation programs for females with PFPS.⁶⁸ Adding a core muscle-strengthening program to the conventional physical therapy management can help improve pain and dynamic balance in female patients with patellofemoral pain syndrome.⁶⁹ DeHaven *et al.*¹⁰ reported that 89% of athletes were able to return to athletic activity after a treatment program that consisted of symptomatic control, a progressive resistance program of isotonic hamstrings and isometric quadriceps exercises, a graduated running program and a maintenance program. Reduction of loading to the patellofemoral joint and surrounding soft tissues by limiting exercise is primary to reducing pain. Substitute activities such as bicycling, swimming, or elliptical should be encouraged. Icing can be beneficial but heat is generally not recommended.

Weakness of the hip musculature may be a risk factor for PFPS; therefore, a proximal strengthening program is recommended. In a study by Earl *et al.*,⁷⁰ nineteen females with PFPS participated in an eight-week program aimed at strengthening the hip and core musculature and improving dynamic malalignment. They reported significant improvements in pain, functional ability, lateral core endurance, hip abduction and hip external rotation strength. They also observed a significant decrease in knee abduction moment during running. These results suggest that an exercise plan that focuses on strengthening and improving neuromuscular control of the hip and core muscles produces positive results in female

Diagnosis

PFPS is a common cause for anterior knee pain and mainly affects young women without any structural changes or significant pathological changes in the articular cartilage. Therefore, PFPS is often known as a diagnosis of exclusion.⁶⁶ Patients with PFPS often describe pain behind, underneath, or around the patella.¹ The symptoms are usually gradual and pain in the anterior knee is the primary symptom of PFPS, although some patients also report instability and crepitation of the patellofemoral joint, specifically during loading of the joint and palpation of the patella. The pain increases after prolonged sitting, squatting, kneeling and stair climbing. PFPS is defined as anterior knee pain or retropatellar pain after at least two of these activities: ascending and descending stairs, hopping, jogging, prolonged sitting, kneeling and squatting.⁶⁷ PFPS also excludes peripatellar tendonitis or bursitis, plica syndromes,

Treatment approaches

Most treatment approaches for PFPS are conservative and surgical interventions are much less common (Table 3). There are a wide variety of treatment programs for PFPS but key components involve increasing strength, flexibility, proprioception, endurance, function training and gradual progression.⁸ A multimodal nonoperative therapy with short-term use of NSAIDs, medially directed tape, and complex exercise programs with the inclusion of the core, lower extremity, and hip and trunk muscles has been shown to be the best course of treatment.⁸

Non-surgical interventions

Females with PFPS had lower eccentric hip abduction and adduction peak torque and higher eccentric adduction to abduction torque ratios when compared with controls. Thus, clinicians should consider eccentric hip abduction strengthening exercises when

Table 1. Patellofemoral pain syndrome (PFPS): summary.

Definition of PFPS	1) Retropatellar pain during stairs, hopping/jogging, prolonged sitting, kneeling, squatting. 2) Negative findings on examination of knee ligament, menisci, bursa, synovial plica. 3) Pain on palpation of patellar facets, femoral condyles.
Incidence/Prevalence	1) Females are twice as likely to develop PFPS compared to males. 2) 70% of cases are between the ages of 16 and 25

Table 2. Reasons for increased susceptibility of patellofemoral pain syndrome in females.

Increase static	q-angle
Increase dynamic	Knee valgus angle; hip internal rotation angle; hip abduction moment; knee valgus moment
Decrease dynamic	Knee flexion angle
Weaker strength of	Quadriceps; hip external rotation; hip extension; hip abducto

Table 3. Patellofemoral pain syndrome treatment options.

Surgical	Non-surgical
Lateral Retinacular Release	Relative Rest
Proximal Realignment Procedures	Physical Therapy
Distal Realignment Procedures	Proximal Strengthening
Elevation of Tibial Tubercle	Gait Retraining
Anteromedial Tibial Tubercle Transfer & Elevation	Analgesics
Articular Cartilage Procedures	Bracing
Patellectomy	Patellar Taping

patients, improves the strength of the hip and core muscles, and reduces knee abduction moment, all of which are associated with the development of PFPS.⁷⁰

There is a large amount of evidence that PFPS is at least partially due to faulty mechanics of the lower extremity.⁷¹ Over time, repetitive exposure to motions such as increased hip adduction and femoral internal rotation may damage or overload the cartilage in the knee joint, which leads to the chronic pain of PFPS.⁷¹ The goal of gait retraining involves adopting new gait patterns and it can be a successful therapy for reducing pain and improving function in PFPS patients as well as long-term improvements.

Although nonsteroidal anti-inflammatory drugs (NSAIDs) are commonly prescribed for patients with PFPS, there is little evidence supporting their effectiveness.⁷² NSAIDs or acetaminophen may be considered for patients whose symptoms cannot be reduced by icing. Furthermore, a variety of braces, sleeves, and straps have been used in the treatment of PFPS. Although bracing alone may provide some symptomatic relief, studies have not found a benefit when bracing is used in addition to physical therapy.⁷³

The recommended approach for patellar taping described by McConnell⁷⁴ is widely cited in treatments for PFPS. Werner *et al.*⁷⁵ found that patients who had patellar hypermobility were able to increase their knee extensor torque by taping. In contrast, Cerny *et al.*⁷⁶ reported that the ratio of vastus medialis obliquus to vastus lateralis as measured by electromyographic activity was not improved with patellar taping. Hence further research is needed to determine whether patellar taping is beneficial in the treatment of PFPS.

Operative intervention

Although surgical interventions are typically not performed due to a wide range of effective conservative treatments available for PFPS, there are a number of surgical procedures that can be performed. Most of these surgical interventions aim at treating malalignment or injured cartilage. Surgical consultation for PFPS may be considered for those patients whose symptoms persist despite completing 6-12 months of conservative management with both formal and home exercises and rehabilitation.

Lateral retinacular release is performed when there is lateral compression syndrome with tenderness and tightness of the lateral retinaculum which is combined with lateral patellar tilt.⁶⁷ Fulkerson *et al.* used CT images to compare patella tilt in patients before and after the lateral release and

found that the lateral release effectively reduces abnormal patella tilt.⁷⁷ Fabbriani *et al.*⁷⁸ found 71% of all patients with patellofemoral pain and presence of patella tilt had satisfactory outcomes after the lateral release procedure. There is a fine balance between too little of a release that will cause persistent pain or too much release that may result in medial patella instability.⁷⁹ Additionally, this procedure is not recommended for very young patients, those with advanced patellofemoral osteoarthritis or patients with normal patellar tracking or patellar tilt.

Proximal realignment procedures are rarely used but are indicated for skeletally immature patients with a history of recurrent dislocations, patients with an increased congruence angle and patients with dysplastic femoral trochlea and poor medial patellar support of the vastus medialis obliquus muscle which leads to recurrent patellar subluxations or dislocations.⁸⁰

Distal realignment or tibial tubercle osteotomy procedures are generally performed on patients with recurrent patellar dislocation or subluxation. Indications for distal realignment procedures include persistent patellofemoral pain coupled with excessive patellar tilt, subluxation or increased congruence angle, as well as lateral facet osteoarthritis in the setting of increased distance between the Tibial Tubercle and the Trochlea Groove (TT-TG). Less than 10% of all patients with PFPS will need a distal realignment procedure. There are several methods for distal realignment, with the most common one being the Fulkerson osteotomy. A 5 to 7 cm bone pedicle is osteotomized at the distal tibial tubercle and the pedicle is moved both anteriorly and medially. The amount of anteriorization versus medialization is dependent on the steepness of the osteotomy cut. The other classic distal realignment also includes the Elmslie-Trillat procedure and Hauser procedure which involves medial translation of the distal tibial tubercle without anterior translation. The Maquet procedure involves anterior translation of the tibial tubercle without medialization to decrease patellofemoral contact forces. Concomitant articular cartilage procedures may also be indicated and include open or arthroscopic shaving of the patella, local excision of defects with drilling of the subchondral bone, and transplantation of autologous chondrocytes or osteochondral allograft procedures.⁸¹ Indications for these procedures are dependent on the status of the cartilage at the time of surgery and is beyond the scope of this review article.

Conclusions

PFPS is one of the most common knee complaints in young healthy female athletes. However, both researchers and clinicians struggle to understand the factors that underlie PFPS because PFPS is often multifactorial in nature and may vary from patient to patient. From a review of the current literature, it is clear that there needs to be further research on PFPS in order to better understand the complex etiology of this disorder in both males and females. Females with patellofemoral pain syndrome demonstrate a decrease in abduction, external rotation and extension strength of the affected side compared with healthy controls.⁴⁸ Due to the complex nature of PFPS, a multitude of treatment approaches have been suggested. However, there is not yet a clear consensus among clinicians regarding the optimal treatment of PFPS. Factors contributing to PFPS include lower extremity malalignment, patellofemoral tilt or balance, muscle imbalance, and soft tissue or cartilage abnormalities. Optimizing the muscle balance between the vastus medialis and lateralis around the patella with formal and home directed therapy should be the first line of treatment in patients presenting with PFPS. Surgery should be reserved for patients with persistent knee pain with defined lesions within the knee, abnormal tilt, and malalignment despite trying all avenues of conservative treatment options.

References

1. Rixe JA, Glick JE, Brady J, Olympia RP. A review of the management of patellofemoral pain syndrome. *Phys Sportsmed* 2013;41:19-28.
2. Taunton JE, Ryan MB, Clement DB, *et al.* A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med* 2002;36:95-101.
3. Tecklenburg K, Dejour D, Hoser C, Fink C. Bony and cartilaginous anatomy of the patellofemoral joint. *Knee Surg Sports Traumatol Arthrosc* 2006;14:235-40.
4. Horton MG, Hall TL. Quadriceps femoris muscle angle: normal values and relationships with gender and selected skeletal measures. *Phys Ther* 1989;69:897-901.
5. Devereaux MD, Lachmann SM. Patello-femoral arthralgia in athletes attending a Sports Injury Clinic. *Br J Sports Med* 1984;18:18-21.
6. Collado H, Fredericson M. Patellofemoral pain syndrome. *Clin*

- Sports Med 2010;29:379-98.
7. Moore K, Dalley II A, Agur A. Clinically Oriented Anatomy. 7th ed: Wolters Kluwer Health; 2014.
 8. Thomee R, Augustsson J, Karlsson J. Patellofemoral pain syndrome: a review of current issues. Sports Med 1999;28:245-62.
 9. Decker MJ, Torry MR, Wyland DJ, et al. Gender differences in lower extremity kinematics, kinetics and energy absorption during landing. Clin Biomech (Bristol, Avon) 2003;18:662-9.
 10. Dehaven KE, Dolan WA, Mayer PJ. Chondromalacia patellae in athletes. Clinical presentation and conservative management. Am J Sports Med 1979; 7:5-11.
 11. Malinzak RA, Colby SM, Kirkendall DT, et al. A comparison of knee joint motion patterns between men and women in selected athletic tasks. Clin Biomech (Bristol, Avon) 2001;16:438-45.
 12. Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. Med Sci Sports Exerc 2003;35: 1745-50.
 13. Baquie P, Brukner P. Injuries presenting to an Australian sports medicine centre: a 12-month study. Clin J Sport Med 1997;7:28-31.
 14. Pollard CD, Sigward SM, Ota S, et al. The influence of in-season injury prevention training on lower-extremity kinematics during landing in female soccer players. Clin J Sport Med 2006; 16:223-7.
 15. Mullaney MJ, Fukunaga T. Current concepts and treatment of patellofemoral compressive issues. Int J Sports Phys Ther 2016;11:891-902.
 16. Powers CM, Bolgla LA, Callaghan MJ, et al. Patellofemoral pain: proximal, distal, and local factors, 2nd International Research Retreat. J Orthop Sports Phys Ther 2012;42:A1-54.
 17. Oakes JL, McCandless P, Selfe J. Exploration of the current evidence base for the incidence and prevalence of patellofemoral pain syndrome. Phys apy Rev 2009;16:382-87.
 18. Thomee R, Renstrom P, Karlsson J, Grimby G. Patellofemoral pain syndrome in young women. I. A clinical analysis of alignment, pain parameters, common symptoms and functional activity level. Scand J Med Sci Sports 1995;5:237-44.
 19. Boling M, Padua D, Marshall S, et al. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. Scand J Med Sci Sports 2010; 20:725-30.
 20. Witvrouw E, Werner S, Mikkelsen C, et al. Clinical classification of patellofemoral pain syndrome: guidelines for non-operative treatment. Knee Surg Sports Traumatol Arthrosc 2005; 13:122-30.
 21. Petersen W, Ellermann A, Gosele-Koppenburg A, et al. Patellofemoral pain syndrome. Knee Surg Sports Traumatol Arthrosc 2014;22:2264-74.
 22. Huberti HH, Hayes WC. Patellofemoral contact pressures. The influence of q-angle and tendofemoral contact. J Bone Joint Surg Am 1984;66:715-24.
 23. Heino Brechter J, Powers CM. Patellofemoral stress during walking in persons with and without patellofemoral pain. Med Sci Sports Exerc 2002;34:1582-93.
 24. Sheehan FT, Derasari A, Fine KM, et al. Q-angle and J-sign: indicative of mal-tracking subgroups in patellofemoral pain. Clin Orthop Relat Res 2010;468:266-75.
 25. Willson JD, Davis IS. Utility of the frontal plane projection angle in females with patellofemoral pain. J Orthop Sports Phys Ther 2008;38:606-15.
 26. Dupuy DE, Hangen DH, Zachazewski JE, et al. Kinematic CT of the patellofemoral joint. AJR Am J Roentgenol 1997;169:211-5.
 27. Adirim TA, Cheng TL. Overview of injuries in the young athlete. Sports Med 2003;33:75-81.
 28. Connolly KD, Ronsky JL, Westover LM, et al. Differences in patellofemoral contact mechanics associated with patellofemoral pain syndrome. J Biomech 2009;42:2802-7.
 29. Farrokhi S, Colletti PM, Powers CM. Differences in patellar cartilage thickness, transverse relaxation time, and deformational behavior: a comparison of young women with and without patellofemoral pain. Am J Sports Med 2011;39:384-91.
 30. Keet JHL, Gray J, Harley Y, Lambert MI. The effect of medial patellar taping on pain, strength and neuromuscular recruitment in subjects with and without patellofemoral pain. Physiotherapy 2007;93:45-52.
 31. Shin SR, Schepsis AA, Murakami A, Edgar CM. Quantification Of Trochlear Dysplasia Via Computed Tomography: Assessment of Morphology Difference Between Control and Chronic Patellofemoral Instability Patients. Orthop J Sports Med 2014;2.
 32. Lin F, Wilson NA, Makhous M, et al. In vivo patellar tracking induced by individual quadriceps components in individuals with patellofemoral pain. J Biomech 2010;43:235-41.
 33. Goh JC, Lee PY, Bose K. A cadaver study of the function of the oblique part of vastus medialis. J Bone Joint Surg Br 1995;77:225-31.
 34. Pattyn E, Verdonk P, Steyaert A, et al. Vastus medialis obliquus atrophy: does it exist in patellofemoral pain syndrome? Am J Sports Med 2011; 39:1450-5.
 35. Cross TM, Gibbs N, Houang MT, Cameron M. Acute quadriceps muscle strains: magnetic resonance imaging features and prognosis. Am J Sports Med 2004;32:710-9.
 36. Callaghan MJ, Oldham JA. Quadriceps atrophy: to what extent does it exist in patellofemoral pain syndrome? Br J Sports Med 2004;38:295-9.
 37. Milgrom C, Finestone A, Eldad A, Shlamkovitch N. Patellofemoral pain caused by overactivity. A prospective study of risk factors in infantry recruits. J Bone Joint Surg Am 1991;73:1041-3.
 38. Herbst KA, Barber Foss KD, Fader L, et al. Hip strength is greater in athletes who subsequently develop patellofemoral pain. Am J Sports Med 2015;43:2747-52.
 39. Kaya D, Citaker S, Kerimoglu U, et al. Women with patellofemoral pain syndrome have quadriceps femoris volume and strength deficiency. Knee Surg Sports Traumatol Arthrosc 2011; 19:242-7.
 40. Lephart SM, Ferris CM, Riemann BL, et al. Gender differences in strength and lower extremity kinematics during landing. Clin Orthop Relat Res 2002:162-9.
 41. Besier TF, Fredericson M, Gold GE, et al. Knee muscle forces during walking and running in patellofemoral pain patients and pain-free controls. J Biomech 2009;42:898-905.
 42. Hehne HJ. Biomechanics of the patellofemoral joint and its clinical relevance. Clin Orthop Relat Res 1990:73-85.
 43. Jonsson P, Alfredson H. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study. Br J Sports Med 2005; 39:847-50.
 44. Hyong IH. Effects of squats accompanied by hip joint adduction on the selective activity of the vastus medialis oblique. J Phys Ther Sci 2015;27:1979-81.
 45. Jan MH, Lin DH, Lin JJ, et al.

- Differences in sonographic characteristics of the vastus medialis obliquus between patients with patellofemoral pain syndrome and healthy adults. *Am J Sports Med* 2009;37:1743-9.
46. Van Tiggelen D, Cowan S, Coorevits P, et al. Delayed vastus medialis obliquus to vastus lateralis onset timing contributes to the development of patellofemoral pain in previously healthy men: a prospective study. *Am J Sports Med* 2009;37:1099-105.
 47. Akkurt E, Salli A, Ozerbil OM, Ugurlu H. The effect of isokenetic exercise on symptoms, functional status and EMC activation onset time of the vastus medialis oblique and vastus lateralis in female patients with patellofemoral pain syndrom. *Isokinet Exerc Sci* 2010;18:157-61.
 48. Karst GM, Willett GM. Onset timing of electromyographic activity in the vastus medialis oblique and vastus lateralis muscles in subjects with and without patellofemoral pain syndrome. *Phys Ther* 1995;75:813-23.
 49. Chester R, Smith TO, Sweeting D, et al. The relative timing of VMO and VL in the aetiology of anterior knee pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2008;9:64.
 50. Ireland ML, Willson JD, Ballantyne BT, Davis IM. Hip strength in females with and without patellofemoral pain. *J Orthop Sports Phys Ther* 2003;33:671-6.
 51. Robinson RL, Nee RJ. Analysis of hip strength in females seeking physical therapy treatment for unilateral patellofemoral pain syndrome. *J Orthop Sports Phys Ther* 2007;37:232-8.
 52. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther* 2009;39:12-9.
 53. Willson JD, Kernozek TW, Arndt RL, et al. Gluteal muscle activation during running in females with and without patellofemoral pain syndrome. *Clin Biomech (Bristol, Avon)* 2011;26:735-40.
 54. Fairbank JC, Pynsent PB, van Poortvliet JA, Phillips H. Mechanical factors in the incidence of knee pain in adolescents and young adults. *J Bone Joint Surg Br* 1984;66:685-93.
 55. Wilson T. The measurement of patellar alignment in patellofemoral pain syndrome: are we confusing assumptions with evidence? *J Orthop Sports Phys Ther* 2007;37:330-41.
 56. Arendt EA. Musculoskeletal injuries of the knee: are females at greater risk? *Minn Med* 2007;90:38-40.
 57. Tenan MS, Peng YL, Hackney AC, Griffin L. Menstrual cycle mediates vastus medialis and vastus medialis oblique muscle activity. *Med Sci Sports Exerc* 2013;45:2151-7.
 58. Cibulka MT, Threlkeld-Watkins J. Patellofemoral pain and asymmetrical hip rotation. *Phys Ther* 2005;85:1201-7.
 59. Prins MR, van der Wurff P. Females with patellofemoral pain syndrome have weak hip muscles: a systematic review. *Aust J Physiother* 2009;55:9-15.
 60. James SL, Bates BT, Osternig LR. Injuries to runners. *Am J Sports Med* 1978;6:40-50.
 61. Hartel MJ, Petersik A, Schmidt A, et al. Determination of Femoral neck angle and torsion angle utilizing a novel three-dimensional modeling and analytical technology based on CT datasets. *PLoS One* 2016;11:e0149480.
 62. Dye SF. Therapeutic implications of a tissue homeostasis approach to patellofemoral pain. *Sports Med Arthrosc Rev* 2001;9:306-11.
 63. Collins N, Crossley K, Beller E, et al. Foot orthoses and physiotherapy in the treatment of patellofemoral pain syndrome: randomised clinical trial. *Br J Sports Med* 2009;43:169-71.
 64. Loudon JK, Wiesner D, Goist-Foley HL, et al. Intrarater reliability of functional performance tests for subjects with patellofemoral pain syndrome. *J Athl Train* 2002;37:256-61.
 65. Casey E, Rho M, Press J. Sex differences in sports medicine. *Demos medical*. 2016.
 66. Al-Hakim W, Jaiswal PK, Khan W, Johnstone D. The non-operative treatment of anterior knee pain. *Open Orthop J* 2012;6:320-6.
 67. Noehren B, Pohl MB, Sanchez Z, et al. Proximal and distal kinematics in female runners with patellofemoral pain. *Clin Biomech (Bristol, Avon)* 2012;27:366-71.
 68. Baldon Rde M, Nakagawa TH, Muniz TB, et al. Eccentric hip muscle function in females with and without patellofemoral pain syndrome. *J Athl Train* 2009;44:490-6.
 69. Chevidikunnan MF, Al Saif A, Gaowgzeh RA, Mamdouh KA. Effectiveness of core muscle strengthening for improving pain and dynamic balance among female patients with patellofemoral pain syndrome. *J Phys Ther Sci* 2016;28:1518-23.
 70. Earl JE, Hoch AZ. A proximal strengthening program improves pain, function, and biomechanics in women with patellofemoral pain syndrome. *Am J Sports Med* 2011;39:154-63.
 71. Noehren B, Scholz J, Davis I. The effect of real-time gait retraining on hip kinematics, pain and function in subjects with patellofemoral pain syndrome. *Br J Sports Med* 2011;45:691-6.
 72. Heintjes E, Berger MY, Bierma-Zeinstra SM, et al. Pharmacotherapy for patellofemoral pain syndrome. *Cochrane Database Syst Rev* 2004; CD003470.
 73. Finestone A, Radin EL, Lev B, et al. Treatment of overuse patellofemoral pain. Prospective randomized controlled clinical trial in a military setting. *Clin Orthop Relat Res* 1993:208-10.
 74. McConnell J. The management of chondromalacia patellae: a long term solution. *Aust J Physiother* 1986;32:215-23.
 75. Werner S, Knutsson E, Eriksson E. Effect of taping the patella on concentric and eccentric torque and EMG of knee extensor and flexor muscles in patients with patellofemoral pain syndrome. *Knee Surg Sports Traumatol Arthrosc* 1993;1:169-77.
 76. Cerny K. Vastus medialis oblique/vastus lateralis muscle activity ratios for selected exercises in persons with and without patellofemoral pain syndrome. *Phys Ther* 1995;75:672-83.
 77. Fulkerson JP, Schutzer SF, Ramsby GR, Bernstein RA. Computerized tomography of the patellofemoral joint before and after lateral release or realignment. *Arthroscopy* 1987;3:19-24.
 78. Fabbriani C, Panni AS, Delcogliano A. Role of arthroscopic lateral release in the treatment of patellofemoral disorders. *Arthroscopy* 1992;8:531-6.
 79. Hughston JC, Deese M. Medial subluxation of the patella as a complication of lateral retinacular release. *Am J Sports Med* 1988;16:383-8.
 80. Weber AE, Nathani A, Dines JS, et al. An Algorithmic approach to the management of recurrent lateral patellar dislocation. *J Bone Joint Surg Am* 2016; 98:417-27.
 81. Parada SA, Eichinger JK, Dumont GD, et al. Comparison of Glenoid version and posterior humeral subluxation in patients with and without posterior shoulder instability. *Arthroscopy* 2017; 33:254-60.