

## Xinning Li, M.D.

Associate Professor  
Sports Medicine and Shoulder Surgery  
Boston University School of Medicine – Boston Medical Center  
Boston University Athletics – Team Physician  
725 Albany Street – 4<sup>th</sup> Floor  
Boston, MA 02118



(617) 638-5633 (Telephone – Shapiro Office)  
(617) 358-3400 (Telephone – Ryan Center)  
(617) 414-5226 (Fax)  
[www.tigerortho.com](http://www.tigerortho.com)



Patient Name:

Date:

Date of surgery:

Visit per week:

### ARTHROSCOPIC KNEE MENISECTOMY PROCEDURE

#### PHYSICAL THERAPY PRESCRIPTION

##### Phase I: Day 1 to 2 Weeks

##### Brace

- Knee immobilizer for comfort only. Please discontinue within one to two weeks after surgery
- Over the counter TED hose can be worn until ambulation has returned to normal pre-surgery level or two weeks
- Crutches as needed for ambulation in the first 2 weeks: weight bearing as tolerated

##### Goals

Decrease inflammation and swelling

- Restore knee range of motion (0 to 115 degrees, minimum 0 degrees extension to 90 degrees flexion before Phase II)
- Re-establish quadriceps muscle activity/re-education
- Weight bearing as tolerated with use of crutches until normal gait is achieved

Note: Return to office for post-op appointment with my Physician Assistant around 7 to 10 days after surgery. Please make sure you have an appointment at the time of surgery.

##### Knee Motion

Passive Range of Motion:

- Prone hang or heel prop (with or without ice)

Active Range of Motion:

- Ankle pumps
- Quadriceps sets
- Straight leg raises
- Hip abduction, adduction, extension
- Gluteal sets
- Heel slides
- Stationary bike (no resistance, for range of motion only)

Strengthening:

- Partial squats (complete with supervision of physical therapist or certified athletic trainer)

Stretching:

- Active assistive range of motion stretching (quadriceps, gastroc, soleus and hamstring)

Modalities

- Neuromuscular electrical stimulation if patient has quadriceps shut down. Thirty minutes, two times a day, seven days a week.

## **Phase II: Weeks 2 to 8**

**Brace:** None

### **Goals**

- Restore and improve muscular strength and endurance
- Gradual return to functional activities
- Restore normal gait
- Improve balance and proprioception
- Re-establish full, pain free range of motion

1. Week two, 0 to 125 degrees
2. Week three, 0 to 135 degrees
3. Week four, 0 to 145 degrees

Cardiovascular

- Stairmaster, elliptical, treadmill (no running) and bike
- Aquatic therapy: running in pool or on AlterG® Anti-Gravity Treadmill® four to six weeks (incision needs to be fully closed)

### **Knee Motion:**

Passive Range of Motion:

- Continue all from Phase I

Active Range of Motion:

- Continue all from Phase I

Strengthening:

- Toe raises and calf raises
- Hamstring curls

- Lunges – front and lateral
- Leg press (not past 90 degrees)
- Step-up, step down, lateral step-up
- Four – way hip: standing
- Knee extension exercise: 90 to 40 degrees (do not use machine)
- TKE
- Closed kinetic chain

Stretching:

- Continue all from Phase I

Proprioception

- Proprioceptive and balance training

Modalities

- Neuromuscular electrical stimulation if patient has quadriceps shut down. Thirty minutes,

two times a day, seven days a week.

### **Phase III: 8 Weeks to 12 Weeks**

#### **Goals**

- Enhance muscular strength and endurance
- Maintain full range of motion

#### **Knee Motion:**

Full flexion and extension

Functional: Begin

- Begin plyometrics/vertical jumps
- Initiate running program
- Agility drills progression
- Forward/backwards running, cutting, figure eight and carioca
- Sports specific drills/activities (begin with non-contact)

**Note:** Monitor patellofemoral joint pain and avoid activities that may aggravate it.

**Signature:** \_\_\_\_\_

Xinning Li, M.D.  
Associate Professor  
Boston University School of Medicine  
[www.tigerortho.com](http://www.tigerortho.com)

