

Xinning Li, M.D.

Professor of Orthopaedic Surgery
Sports Medicine and Shoulder Surgery
Boston University School of Medicine – Boston Medical Center
Boston University Sports Medicine Fellowship – Director
Boston University Athletics – Team Physician



725 Albany Street – 4th Floor
Boston, MA 02118
(617) 638-5633 (Telephone – Shapiro Office)
(617) 358-3400 (Telephone – Ryan Center)
(617) 414-5226 (Fax)
www.tigerortho.com



RIGHT or LEFT KNEE MCL RECONSTRUCTION W/ ALLOGRAFT

Week 1-2 (Ankle pumps every hour)

- Post – op brace locked to maintain full extension for 2 weeks
- Quad sets & SLR (Brace on) with no lag
- WBAT with crutches
- Ice or Cryocuff Unit on knee for 20 – 30 minutes every hour
- Passive ROM exercises: Limits: 0 to 40 degrees.
- NO Hip adductor strengthening

Week 3-4 (ROM 0-75deg, WBAT)

- Supervised PT –2- 3 times a week (may need to adjust based on insurance)
- Continue SLR's in brace with foot straight up, quad isometric sets, ankle pumps
- WBAT with brace opened full ROM.
- Patellar mobilization exercises and edema control.
- Brace opened for full ROM for ambulation and sleeping.
- May remove brace for HEP and NO Hip adductor strengthening.

Week 5 (ROM as tolerated, WBAT)

- Continue with above exercises/ice treatments.
- Advance ROM as tolerated with no limits with brace on.
- Stationary bike for range of motion (short crank or high seat, no resistance) Ok to remove brace for bike here.
- Weight bearing as tolerated with knee in flexed position, start to transition out of the brace.

- Perform scar message aggressively
- Progressive SLR program for quad strength with brace on – start with 1 lb, progress 1-2 lbs per week
- Hamstring and hip PREs
- Seated leg extension (90 to 40degrees) against gravity with no weight
- NO side lying Hip adductor strengthening

Week 6 (WBAT)

- Continue all exercises
- WBAT on the operative knee and transition out of the brace
- Flexion exercises seated AAROM
- AAROM (using good leg to assist) exercises (4-5x/ day) with brace on
- Continue ROM stretching and overpressure into extension
- SLR's – with brace on
- NO side lying Hip adductor strengthening
- Leg press 0-70 arc of motion

Week 7 (WBAT)

- Continue above exercises
- Out of the Brace and Ambulate without crutches.
- Hamstring and calf stretching
- Self ROM 4-5x/day using other leg to provide ROM
- Advance ROM as tolerated – no limits, may remove brace for ROM
- Regular stationary bike if Flexion > 115
- Heel raises with brace on
- Hip strengthening No side lying hip adduction

Week 8

- Continue above exercises
- Mini squats (0-60 degrees)
- 4 inch step ups
- Isotonic leg press (0 – 90 degrees)
- Lateral step out with therabands
- Hip strengthening

Week 9

Advance ROM, Goal: 0 to 125 degrees, walking with no limp

- Add ball squats
- Initiate retro treadmill with 3% incline (for quad control)
- Increase resistance on stationary bike
- Mini-squats and weight shifts
- Sportcord (bungee) walking
- 8 inch step ups
- 4 inch step downs

Week 10

- Begin resistance for open chain knee extension
- Swimming allowed, flutter kick only
- Bike outdoors, level surfaces only
- Progress balance and board throws
- Plyometric leg press
- 6-8 inch step downs
- Start slide board
- Jump down's (double stance landing)
- Progress to light running program and light sport specific drills if:
 - Quad strength > 75% contralateral side
 - Active ROM 0 to >125 degrees
 - Functional hop test >70% contralateral side
 - Swelling < 1cm at joint line
 - No pain
 - Demonstrates good control on step down

Week 11-22

- Stair master machine
- If full ROM, quad strength > 80% contralateral side, functional hop test >85% contralateral side, satisfactory clinical exam:
 - Progress to home program for running. Progress to hops, jumps, cuts and sports specific drills. Begin to wean from supervised therapy.

4 – 5 months

- Criteria to return to sports
 - Full Active ROM
 - Quadriceps >90% contralateral side
 - Satisfactory clinical exam
 - Functional hop test > 90% contralateral side
 - Completion of ACL running program

One Year

- Doctor visit

This is strictly an outline of most of the major exercises that we would like to incorporate into the MCL rehabilitation. Not all exercises need to be done. Two main goals are that appropriate progress is made on a weekly basis, and that communication exists between patient, therapist and doctor.

Physician's Signature: _____

Xinning Li, M.D.
Professor of Orthopaedic Surgery
Boston University School of Medicine
www.tigerortho.com

