

Xinning Li, M.D.

Associate Professor
Sports Medicine and Shoulder Surgery
Boston University School of Medicine – Boston Medical Center
Boston University Athletics – Team Physician
725 Albany Street – 4th Floor
Boston, MA 02118



(617) 638-5633 (Telephone – Shapiro Office)
(617) 358-3400 (Telephone – Ryan Center)
(617) 414-5226 (Fax)
www.tigerortho.com



Patient Name:

Date:

Date of surgery:

Visit per week:

PHYSICAL THERAPY PRESCRIPTION:

REVERSE SHOULDER ARTHROPLASTY

Weeks 2-6: Phase I

Sling Immobilizer: Please Transition out of the sling between 2 to 3 weeks after surgery (Sleep in sofa recliner with your sling for the first 2 weeks) for Primary Reverse Shoulder Replacement or 5 to 6 weeks for Fracture and Revision Reverse Shoulder Replacement.

- 2 Weeks
- 6 Weeks

Activities of Daily Living: You may use your arm and hand on the operative shoulder for your ADL activates 2 weeks after surgery. Pendulums are also encouraged 1 week after surgery.

Exercises: Passive ER to 20 degrees and extension to neutral, pendulums daily
Passive supine FF **in scapular plane** to 90 (**NO aggressive stretching and NO PAIN**)
AROM wrist/elbow
Submax (50%) pain free deltoid isometrics in neutral
Scapular “pinches”
Modalities as needed: ice

Advancement Criteria: ER to the above set criteria
FF in scapular plane to 90
Minimal pain and inflammation with any motion or exercise
Incision is well healed with no drainage or redness

Weeks 6-10: Phase II

Exercises: Passive & Active assisted FF in scapular plane – no limits (wand exercises, pulleys)
Passive & Active assisted ER as tolerated by the patient
Active supine FF in scapular plane
Manual scapular side-lying stabilization exercises
Isometrics: Deltoid in neutral
ER (modified neutral) ROM < 30 deg
IR (modified neutral)
Scapular retraction with elastic bands
Humeral head control exercises:
ER/IR (supine/scapular plane)
Elevation at 100 deg
Modalities as needed

Advancement Criteria: FF > 120
ER > 30
Good humeral head control
Minimal to no pain with ADLs

Weeks 10-16: Phase III

Exercises: Progress ROM as tolerated
AAROM for full FF and ER
AAROM for IR – no limits
Flexibility exercises: towel stretch, posterior capsule stretch
IR/ER/FF isotonic strengthening
Scapular stabilization
Rhythmic stabilization
PREs for scapula, elbow (biceps/triceps)
Forward flexion in scapular plane
Progressive resistive equipment: row, chest press (light weight)
Modalities as needed

Advancement Criteria: Muscle strength 4/5
Passive FF > 120, ER >30
Restore normal scapulohumeral rhythm <90 deg elevation
Minimal pain and inflammation

Weeks 16-22: Phase IV

Exercises: Access and address any remaining deficits in ROM, flexibility, strength
Active, active-assisted, and passive ROM exercises
Flexibility exercises: towel stretch (IR), posterior capsule stretch
Progressive resistive strengthening:
Dumbbells
Progressive resistive equipment
Elastic band IR/ER (modified neutral)

Rhythmic stabilization
Modalities as needed
Individualize program to meet specific needs of patient

Discharge Criteria: Maximize ROM
Full independent ADLs
Normal scapulohumeral rhythm >100deg elevation

* Please Send Progress Notes *

Signature: _____

Xinning Li, M.D.
Associate Professor
Boston University School of Medicine
www.tigerortho.com

